

## Routine Schedule

Age*	Vaccine	Primary Doses	Booster Dose†
6 months– 4 years	Pfizer– Infant/Toddler	1st Dose → 3-8 weeks <sup>^</sup> → 2nd Dose → ≥8 weeks → 3rd Dose	
6 months– 5 years	Moderna– Infant/Toddler	1st Dose → 4-8 weeks <sup>^</sup> → 2nd Dose	
5–11 years	Pfizer– Pediatric	1st Dose → 3-8 weeks <sup>^</sup> → 2nd Dose	<p><b>Bivalent Booster†</b></p> <ul style="list-style-type: none"> <li>● Pfizer: Ages 5-11</li> <li>● Pfizer: Ages 12+</li> <li>● Moderna: Ages 6+</li> </ul> <p>(For people who previously received a monovalent booster dose(s), the bivalent booster is administered at least 2 months after the last monovalent booster dose.)</p>
6–11 years	Moderna– Pediatric	1st Dose → 4-8 weeks <sup>^</sup> → 2nd Dose	
12+ years	Moderna– Adol/Adult	1st Dose → 4-8 weeks <sup>^</sup> → 2nd Dose	
12+ years	Pfizer/ Adol/Adult	1st Dose → 3-8 weeks <sup>^</sup> → 2nd Dose	
12+ years	Novavax	1st Dose → 3-8 weeks <sup>^</sup> → 2nd Dose	
18+ years	Janssen (J&J) Pfizer/Moderna/ Novavax preferred**	1st Dose	

\* See schedules for children in transition from a younger to older age group: [Pfizer](#) | [Moderna](#).

\*\* Although use of mRNA COVID-19 and Novavax vaccines is preferred, the Janssen vaccine may be offered in [some situations](#).

† For people who have not received any booster doses and are unable or unwilling to receive bivalent booster vaccine, the [monovalent Novavax booster may be administered as a single booster dose](#) at least 6 months after completion of the primary series to people 18 years and older.

<sup>^</sup> An [8-week interval](#) may be preferable for some people, especially for males 12-39 years.

View [Interim Clinical Considerations for Use of COVID-19 Vaccines](#) for details. Schedule is subject to change.

## Schedule if Moderately or Severely Immunocompromised

Age*	Vaccine	Primary Doses	Booster Dose†
6 months–4 years	Pfizer–Infant/Toddler	1st Dose → 3 weeks → 2nd Dose → ≥8 weeks → 3rd Dose	<p><b>Bivalent Booster†</b></p> <ul style="list-style-type: none"> <li>● Pfizer: Ages 5-11</li> <li>● Pfizer: Ages 12+</li> <li>● Moderna: Ages 6+</li> </ul> <p>(For people who previously received a monovalent booster dose(s), the bivalent booster is administered at least 2 months after the last monovalent booster dose.)</p>
6 months–5 years	Moderna–Infant/Toddler	1st Dose → 4 weeks → 2nd Dose → ≥4 weeks → 3rd Dose	
5–11 years	Pfizer–Pediatric	1st Dose → 3 weeks → 2nd Dose → ≥4 weeks → 3rd Dose	
6–11 years	Moderna–Pediatric	1st Dose → 4 weeks → 2nd Dose → ≥4 weeks → 3rd Dose	
12+ years	Moderna–Adol/Adult	1st Dose → 4 weeks → 2nd Dose → ≥4 weeks → 3rd Dose	
12+ years	Pfizer/Adol/Adult	1st Dose → 3 weeks → 2nd Dose → ≥4 weeks → 3rd Dose	
12+ years	Novavax	1st Dose → 3 weeks → 2nd Dose	
18+ years	Janssen (J&J) Pfizer/Moderna/Novavax preferred**	1st Dose → 4 weeks → 2nd Dose of Moderna/Pfizer	

\* See schedules for children in transition from a younger to older age group: [Pfizer](#) | [Moderna](#).

\*\* Although use of mRNA COVID-19 and Novavax vaccines is preferred, the Janssen vaccine may be offered in [some situations](#).

† For people who have not received any booster doses and are unable or unwilling to receive bivalent booster vaccine, the [monovalent Novavax booster may be administered as a single booster dose](#) at least 6 months after completion of the primary series to people 18 years and older.

View [Interim Clinical Considerations for Use of COVID-19 Vaccines](#) for details. Schedule is subject to change.